

Marshall et al J. Neurosurg volume 75, November 1991

Diagnostic categories of types of abnormalities visualized on CT scanning*

Category	Definition
Diffuse Injury I (no visible pathology)	no visible intracranial pathology seen on CT scan
Diffuse Injury II	cisterns are present with midline shift 0-5 mm and/or: lesion densities present no high- or mixed-density lesion > 25 cc may include bone fragments and foreign bodies
Diffuse Injury III (swelling)	cisterns compressed or absent with midline shift 0-5 mm, no high- or mixed-density lesion > 25 cc
Diffuse Injury IV (shift)	midline shift > 5 mm, no high- or mixed-density lesion > 25 cc
evacuated mass lesion	any lesion surgically evacuated
nonevacuated mass lesion	high- or mixed-density lesion > 25 cc, not surgically evacuated

Outcome at discharge in relation to intracranial diagnosis*

Outcome at Discharge	Diffuse Injury I		Diffuse Injury II		Diffuse Injury III		Diffuse Injury IV		Evacuated Mass		Nonevacuated Mass		Brain-Stem Injury		Unknown		Totals	
	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%
good	14	27.0	15	8.5	5	3.3	1	3.1	14	5.1	1	2.8	0	0	0	0	50	7.0
moderate	18	34.6	46	26.0	20	13.1	1	3.1	49	17.7	3	8.3	0	0	1	5.9	138	18.5
severe	10	19.2	72	40.7	41	26.8	6	18.8	72	26.1	7	19.4	1	33.3	0	0	209	28.0
vegetative	5	9.6	20	11.3	35	22.9	6	18.8	34	12.3	6	16.7	0	0	0	0	106	14.0
dead	5	9.6	24	13.5	52	34.0	18	56.2	107	38.8	19	52.8	2	66.7	16	94.1	243	32.5
totals	52	100	177	100	153	100	32	100	276	100	36	100	3	100	17	100	746	100

* Outcome classified by the Glasgow Outcome Scale.* For a description of Diffuse Injury 1 to IV, see Table 1.

Score Classification	Essential features
Death	
Persistent vegetative state	Eyes open, sleep-wake cycles. No speech, communication or response to external stimuli.
Severe disability	Lower grade: usually dependent in activities of daily living and institutionalized. May be living at home with a large degree of family and nursing support. Upper grade: may be largely self-caring, but unable to work, even in a sheltered environment, or travel independently.
Moderate disability	Lower grade: able to travel independently, work in a sheltered environment or at lesser level than pre-injury. Upper grade: able to work in a reduced capacity, may have deficits in speech, memory and personality change.
Good recovery	Lower grade: able to participate in social activities outside the home, but to a less extent than pre-injury. Upper grade: resumption of normal life, minor neurological or psychological deficits may persist.

Extended Glasgow Outcome Scale

prognosis after traumatic brain injury [created by Paul Young 14/11/07]

CT findings

general

- predicting outcome after TBI can help guide acute & chronic care & help prepare the family for a typically protracted recovery process
- equally important is that further treatment may be deemed futile & expensive critical care and surgery can thus be reserved for patients who will benefit

predictors

- several clinical and radiological characteristics have proved useful for outcome prediction but they must be used in concert. These criteria are more useful for predicting death or vegetative state than for accurately predicting mild or no dysfunction
- the most powerful outcome predictors are:
 - age
 - initial GCS score
 - pupil size and reaction to light
 - ICP
 - nature & extent of intracranial injuries

age

- old age correlates most consistently with a poor outcome after traumatic brain injury
- traumatic coma data bank study or >700 patients with severe TBI showed that among patients older than 60 years the incidence of death, persistent vegetative state or severe disability was 92%, 86% for those older than 56 and 50% for younger patients
- older patients are more likely to have traumatic intracranial mass lesions & the presence of these insults strongly correlates with poor outcome
- subsequent studies have demonstrated the low probability of a good recovery for patients older than 60 whose initial GCS is <8

GCS

- the second most important predictor of outcome is the initial post-resuscitation GCS score. Among patients with a severe closed head injuries in the traumatic coma data bank study, good outcomes occurred in 4.1% of those with an initial GCS of 3, in 6.3% whose score was 4 and 12.2% whose score was 5

pupils

- unilateral or bilaterally dilated pupils that are unreactive to light usually reflect uncal herniation and significant brainstem compression and damage
- several large clinical studies have found that patients with bilaterally fixed and dilated pupils had a greater than 90% likelihood of death or vegetative state

intracranial hypertension

- intracranial hypertension refractory to medication is associated with a 43% mortality rate and a 0% chance of functional outcome

lesion nature

- subdural haematoma:
 - subdural haematoma is associated with the worst prognosis
 - one study found only 26% of patients with these clots had a functional recovery
 - prognosis is related to how rapidly the clot is evacuated with the best outcomes occurred in patients who have surgery within 2 hours
- extradural haematoma:
 - pose a much lower risk of mortality than SDH because they are not usually associated with underlying cerebral contusion and swelling
 - mortality depends a great deal on time to surgery; untreated lesions can lead to uncal herniation and death
 - mortality increases from 17-65% if an extradural haematoma is not evacuated within 2 hours after the onset of coma
- traumatic subarachnoid:
 - the presence of traumatic subarachnoid haemorrhage is associated with a greater than 50% risk of death; however, many believe that this condition is merely a marker of more severe brain injury and has no direct association with outcome